

INTERNATIONAL CENTRE FOR ECONOMIC RESEARCH



WORKING PAPER SERIES

Fabio Maccheroni and Massimo Marinacci

A STRONG LAW OF LARGE NUMBERS FOR CAPACITIES

Working Paper no. 28/2004
December 2004

**APPLIED MATHEMATICS
WORKING PAPER SERIES**



A strong law of large numbers for capacities*

Fabio Maccheroni^a and Massimo Marinacci^b

^aIstituto di Metodi Quantitativi and IGIER, Università Bocconi

Viale Isonzo 25, 20135 - Milano, ITALY

fabio.maccheroni@unibocconi.it

<http://web.econ.unito.it/gma/fabio.htm>

^bDipartimento di Statistica e Matematica Applicata and ICER, Università di Torino

Piazza Arbarello 8, 10122 - Torino, ITALY

massimo.marinacci@unito.it

<http://web.econ.unito.it/gma/massimo.htm>

June 2004

Abstract

We consider a totally monotone capacity on a Polish space and a sequence of bounded p.i.i.d. random variables. We show that, on a full set, any cluster point of empirical averages lies between the lower and the upper Choquet integrals of the random variables, provided either the random variables or the capacity are continuous.

Keywords and phrases: Capacities; Choquet integral; Strong law of large numbers.

MSC2000: 28A12 Contents, measures, outer measures, capacities; 60F15 Strong theorems.

*We thank Steve Lalley and an anonymous referee of *The Annals of Probability* for helpful suggestions. The financial support of the *Ministero dell'Istruzione, dell'Università e della Ricerca* is gratefully acknowledged.

1 Introduction

In this paper we prove a strong law of large numbers for totally monotone capacities. Specifically, given a totally monotone capacity ν defined on the Borel σ -algebra \mathcal{B} of a Polish space Ω , and a sequence $\{X_n\}_{n \geq 1}$ of bounded, pairwise independent and identically distributed random variables, we show that

$$\nu \left(\left\{ \omega \in \Omega : \int X_1 d\nu \leq \underline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} \leq \overline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} \leq - \int -X_1 d\nu \right\} \right) = 1,$$

provided the X_n s are continuous or simple, or ν is continuous. In this way we extend earlier results of Marinacci [13].

Under different names, totally monotone capacities have been widely studied in both pure and applied mathematics. They have been introduced by Choquet [4] motivated by some problems in potential theory, and in his wake many works have studied them in both potential theory and probability theory (see, e.g., [6] and [8]).

In mathematical statistics and in mathematical economics, totally monotone capacities have been used to represent subjective prior beliefs when the information on which such beliefs are based is not good enough to represent them by a standard additive probability (see, e.g., [7], [11], [20], [15], and [19])

Our result shows that even in a non-additive setting the limit behavior of empirical averages has some noteworthy properties. In particular, we show that eventually empirical averages lie, with probability one, between the lower and upper Choquet integrals associated with the given capacity. This extends to the non-additive setting the classic Kolmogorov limit law, to which our result reduces when ν is additive since in this case lower and upper Choquet integrals coincide.

In a subjective probability perspective, our result says that, while in the additive case a Bayesian decision maker believes that the empirical averages of a sequence of p.i.i.d. random variables tend to a given number, here he just believes that the limit behavior of the empirical averages is confined in a given interval. This reflects a possible lack of confidence in his probability assessments. This interpretation of non-additive limit laws and its relevance in mathematical economics has been recently discussed at length in Epstein and Schneider [9], to which we refer the interested reader for details and references.

2 Preliminaries

Let Ω be a Polish space and \mathcal{B} its Borel σ -algebra. A *random variable* (r.v.) is a (Borel) measurable function $X : \Omega \rightarrow \mathbb{R}$. A *totally monotone capacity* on \mathcal{B} is a set function

$\nu : \mathcal{B} \rightarrow [0, 1]$ such that

$$(c.1) \quad \nu(\emptyset) = 0 \text{ and } \nu(\Omega) = 1,$$

$$(c.2) \quad \nu(A) \leq \nu(B) \text{ for all Borel sets } A \subseteq B,$$

$$(c.3) \quad \nu(B_n) \downarrow \nu(B) \text{ for all sequences of Borel sets } B_n \downarrow B,$$

$$(c.4) \quad \nu(G_n) \uparrow \nu(G) \text{ for all sequences of open sets } G_n \uparrow G,$$

$$(c.5) \quad \nu\left(\bigcup_{j=1}^n B_j\right) \geq \sum_{\emptyset \neq J \subseteq \{1, \dots, n\}} (-1)^{|J|+1} \nu\left(\bigcap_{j \in J} B_j\right) \text{ for every collection } B_1, \dots, B_n \text{ of Borel sets.}$$

A set function $\nu : \mathcal{B} \rightarrow [0, 1]$ such that

$$(c.6) \quad \nu(B_n) \uparrow \nu(\Omega) \text{ for all sequences of Borel sets } B_n \uparrow \Omega,$$

is called *continuous*. A continuous set function $\nu : \mathcal{B} \rightarrow [0, 1]$ is a totally monotone capacity if and only if (c.1), (c.2), and (c.5) hold (see [18] and [14, Theorem 10]).

Let ν be a totally monotone capacity on \mathcal{B} . As in the additive case, we say that the elements of a sequence $\{X_n\}_{n \geq 1}$ of r.v.s are *pairwise independent* with respect to ν if, for each $n, m \geq 1$ and for all open subsets G_n, G_m of \mathbb{R} ,

$$\nu(\{X_n \in G_n, X_m \in G_m\}) = \nu(\{X_n \in G_n\}) \nu(\{X_m \in G_m\});$$

we say that they are *identically distributed* if, for each $n, m \geq 1$ and each open subset G of \mathbb{R} ,

$$\nu(X_n \in G) = \nu(X_m \in G).$$

The *Choquet integral* of a bounded r.v. X with respect to a totally monotone capacity ν is defined by

$$\int X d\nu \equiv \int_0^{+\infty} \nu(\{X > t\}) dt + \int_{-\infty}^0 [\nu(\{X > t\}) - 1] dt.$$

The integrals on the right are Riemann integrals and they are well defined since $\nu(\{X > t\})$ is a monotone function in t . The Choquet integral is positively homogeneous, monotone, and translation invariant (i.e., $\int (X + c) d\nu = \int X d\nu + c$ if c is constant). It reduces to the standard integral when ν is an additive probability measure.

In general, $\int X d\nu \leq - \int -X d\nu$. Equality holds for all r.v.s if and only if ν is additive. The integrals $\int X d\nu$ and $- \int -X d\nu$ are sometimes called *lower* and *upper Choquet integrals*, respectively.

3 The law of large numbers

We can now state our main result.

Theorem 1 *Let ν be a totally monotone capacity on \mathcal{B} , and $\{X_n\}_{n \geq 1}$ a sequence of bounded, pairwise independent and identically distributed random variables. Then*

$$\nu \left(\left\{ \omega \in \Omega : \int X_1 d\nu \leq \underline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} \leq \overline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} \leq - \int -X_1 d\nu \right\} \right) = 1,$$

provided at least one the following two conditions holds:

(i) ν is continuous;

(ii) the random variables X_n are either continuous or simple.

Few remarks are in order. First, as $\nu(B) = 1$ and $A \subseteq B^c$ imply $\nu(A) = 0$, under the assumptions of Theorem 1 we also have

$$\begin{aligned} \nu \left(\left\{ \omega \in \Omega : \underline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} < \int X_1 d\nu \right\} \right) &= 0, \text{ and} \\ \nu \left(\left\{ \omega \in \Omega : \overline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} > - \int -X_1 d\nu \right\} \right) &= 0. \end{aligned}$$

In other words, with zero probability empirical averages will eventually lie outside the interval $[\int X_1 d\nu, - \int -X_1 d\nu]$.

Second, when ν is additive we have $\int X_1 d\nu = - \int -X_1 d\nu$, and so in this case our result reduces to a standard Kolmogorov limit law

$$\nu \left(\left\{ \omega \in \Omega : \lim_n \frac{\sum_{j=1}^n X_j(\omega)}{n} = \int X_1 d\nu \right\} \right) = 1.$$

On the other hand, when ν is not additive in some cases it may happen (see [13]) that

$$\nu \left(\left\{ \omega \in \Omega : \underline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} < \overline{\lim}_n \frac{\sum_{j=1}^n X_j(\omega)}{n} \right\} \right) = 1.$$

Finally, as anticipated, the closest existing theorem is due to [13]. Our result is more general since [13] assumes that Ω is compact, ν is continuous, the r.v.s X_n are continuous, independent, and that they satisfy some further technical conditions. Moreover, the proof we provide is different and much simpler. In fact, here we develop a technique that relies on the relations between totally monotone capacities and correspondences, thus making it possible to use existing laws of large numbers for correspondences. This approach might be useful in establishing further generalizations of limit laws to the framework of capacities. This will be the object of future research, along with the possibility of weakening some of the continuity conditions assumed in Theorem 1.

4 Proof and related material

Denote by \mathcal{K}_Ω (resp., \mathcal{G}_Ω) the class of all nonempty compact subsets (resp., open subsets) of Ω ; for the sake of completeness write \mathcal{B}_Ω instead of \mathcal{B} . If d is a Polish metric on Ω , then \mathcal{K}_Ω is a Polish space when endowed with the Hausdorff metric

$$d_{\mathcal{H}}(K, L) \equiv \max \left(\max_{k \in K} \min_{l \in L} d(k, l), \max_{l \in L} \min_{k \in K} d(l, k) \right).$$

The Borel σ -algebra on \mathcal{K}_Ω is also generated by the class $\{K \in \mathcal{K}_\Omega : K \subseteq G\}_{G \in \mathcal{G}_\Omega}$.

4.1 Measurable correspondences and totally monotone capacities

Let $(I, \mathcal{C}, \lambda)$ be a non-atomic and complete probability space. A (*compact valued*) *correspondence* $F : I \rightrightarrows \Omega$ is a map with domain I and whose values are nonempty compact subsets of Ω . For any $A \subseteq \Omega$ we put

$$F_{-1}(A) \equiv \{s \in I : F(s) \subseteq A\}.$$

A correspondence $F : I \rightrightarrows \Omega$ is *measurable* if $F_{-1}(G) \in \mathcal{C}$ for every $G \in \mathcal{G}_\Omega$. As well known (see, e.g., [12]), the following facts are equivalent:

- F is measurable;
- $F_{-1}(B) \in \mathcal{C}$ for every $B \in \mathcal{B}_\Omega$;
- F is measurable as a function $F : I \rightarrow \mathcal{K}_\Omega$.

When a measurable correspondence F is regarded as a measurable function $F : I \rightarrow \mathcal{K}_\Omega$, we denote by F^{-1} its standard inverse image, that is,

$$F^{-1}(\mathcal{E}) \equiv \{s \in I : F(s) \in \mathcal{E}\} \quad \forall \mathcal{E} \subseteq \mathcal{K}_\Omega,$$

and by $\sigma(F)$ the σ -algebra generated by F , that is,

$$\sigma(F) \equiv \{F^{-1}(\mathcal{D}) : \mathcal{D} \in \mathcal{B}_{\mathcal{K}_\Omega}\}.$$

In the sequel we will need the next lemma, whose standard proof is omitted.

Lemma 1 *Let $F : I \rightrightarrows \Omega$ be a measurable correspondence. Then*

$$\sigma(F) = \sigma(\{F_{-1}(G) : G \in \mathcal{G}_\Omega\})$$

and $\{F_{-1}(G) : G \in \mathcal{G}_\Omega\}$ is a π -class containing I .

A measurable function $f : I \rightarrow \Omega$ induces a *probability distribution* P_f on \mathcal{B}_Ω defined by

$$P_f(B) \equiv \lambda(f^{-1}(B)) \quad \forall B \in \mathcal{B}_\Omega.$$

In a similar way, a measurable correspondence $F : I \rightrightarrows \Omega$ induces a *lower distribution* ν_F on \mathcal{B}_Ω defined by

$$\nu_F(B) \equiv \lambda(F_{-1}(B)) \quad \forall B \in \mathcal{B}_\Omega.$$

The next result, which links totally monotone capacities and lower distributions, is essentially due to Choquet [4] (see also [15], [16], and [3]).

Lemma 2 *A set function $\nu : \mathcal{B}_\Omega \rightarrow [0, 1]$ is a totally monotone capacity if and only if there exists a measurable correspondence $F : I \rightrightarrows \Omega$ such that $\nu = \nu_F$.*

A *measurable selection* of a correspondence $F : I \rightrightarrows \Omega$ is a measurable function $f : I \rightarrow \Omega$ such that $f(s) \in F(s)$ for almost all $s \in I$. The set of all measurable selections of F is denoted by $\text{Sel } F$. The *Aumann integral* (see [2]) of a correspondence $F : I \rightrightarrows \mathbb{R}$ with respect to λ is defined by

$$\int F d\lambda \equiv \left\{ \int f d\lambda : f \in \text{Sel } F \text{ and } f \text{ integrable} \right\}.$$

If $X : \Omega \rightarrow \mathbb{R}$ is continuous or simple, and F is a correspondence, then $(X \circ F)(s) \equiv X(F(s))$ is a correspondence (i.e., $X(F(s)) \in \mathcal{K}_\mathbb{R}$ for all $s \in I$). Moreover, since

$$(X \circ F)_{-1}(A) = F_{-1}(X^{-1}(A)) \quad \forall A \subseteq \mathbb{R},$$

$X \circ F$ is measurable provided X and F are measurable.

Lemma 3 *Let $F : I \rightrightarrows \Omega$ be a measurable correspondence, and $X : \Omega \rightarrow \mathbb{R}$ be either bounded and continuous or simple and measurable. Then,*

$$\int (X \circ F) d\lambda = \left[\int X d\nu_F, - \int -X d\nu_F \right].$$

This is an immediate consequence of [3, Theorem 2].

4.2 Proof of Theorem 1

Suppose first that (ii) holds. By Lemma 2, there exists a measurable correspondence $F : I \rightrightarrows \Omega$ such that $\nu = \nu_F$.

Next we show that the measurable correspondences $\{X_n \circ F\}_{n \geq 1}$ are pairwise independent and identically distributed when regarded as measurable functions $X_n \circ F : I \rightarrow \mathcal{K}_\mathbb{R}$.

Let $n, m \geq 1$ and $G_n, G_m \in \mathcal{G}_{\mathbb{R}}$, then

$$\begin{aligned}
& \lambda((X_n \circ F)_{-1}(G_n) \cap (X_m \circ F)_{-1}(G_m)) \\
&= \lambda(F_{-1}(X_n^{-1}(G_n)) \cap F_{-1}(X_m^{-1}(G_m))) \\
&= \lambda(F_{-1}(X_n^{-1}(G_n) \cap X_m^{-1}(G_m))) \\
&= \nu(X_n^{-1}(G_n) \cap X_m^{-1}(G_m)) \\
&= \nu(X_n^{-1}(G_n)) \nu(X_m^{-1}(G_m)) \\
&= \lambda(F_{-1}(X_n^{-1}(G_n))) \lambda(F_{-1}(X_m^{-1}(G_m))) \\
&= \lambda((X_n \circ F)_{-1}(G_n)) \lambda((X_m \circ F)_{-1}(G_m)).
\end{aligned}$$

This proves pairwise independence, since for all $j = n, m$, $\{(X_j \circ F)_{-1}(G)\}_{G \in \mathcal{G}_{\mathbb{R}}}$ is a π -class containing I and generating the σ -algebra $\sigma(X_j \circ F)$ (see Lemma 1).

Moreover, for each $n, m \geq 1$, and each open subset $G \in \mathcal{G}_{\mathbb{R}}$

$$\begin{aligned}
& \lambda((X_n \circ F)^{-1}(\{K \in \mathcal{K}_{\mathbb{R}} : K \subseteq G\})) \\
&= \lambda(\{X_n \circ F \in \{K \in \mathcal{K}_{\mathbb{R}} : K \subseteq G\}\}) \\
&= \lambda((X_n \circ F)_{-1}(G)) \\
&= \lambda(F_{-1}(X_n^{-1}(G))) \\
&= \nu(X_n^{-1}(G)) \\
&= \nu(X_m^{-1}(G)) \\
&= \lambda((X_m \circ F)^{-1}(\{K \in \mathcal{K}_{\mathbb{R}} : K \subseteq G\})).
\end{aligned}$$

This proves identical distribution since $\{K \in \mathcal{K}_{\mathbb{R}} : K \subseteq G\}_{G \in \mathcal{G}_{\mathbb{R}}}$ is a π -class containing $\mathcal{K}_{\mathbb{R}}$ and generating $\mathcal{B}_{\mathcal{K}_{\mathbb{R}}}$.

Clearly, for each $n \geq 1$ and each $h \in \text{Sel } X_n \circ F$, $\int h d\lambda$ is finite (h is almost surely bounded); moreover, by Lemma 3, $\int X_n \circ F d\lambda \in \mathcal{K}_{\mathbb{R}}$.

In sum, $\{X_n \circ F\}_{n \geq 1}$ are pairwise independent and identically distributed measurable correspondences with $\int X_n \circ F d\lambda \in \mathcal{K}_{\mathbb{R}}$ for all $n \geq 1$. A generalization due to [10] (see also [5] and [17]) of a result of Artstein and Vitale [1], guarantees that

$$\lambda\left(\left\{s \in I : \frac{1}{n} \sum_{j=1}^n X_j(F(s)) \rightarrow \int X_1 \circ F d\lambda\right\}\right) = 1.$$

By Lemma 3,

$$\int X_1 \circ F d\lambda = \left[\int X_1 d\nu, - \int -X_1 d\nu \right].$$

Let $a_n(\omega) = \frac{\sum_{j=1}^n X_j(\omega)}{n}$ and set

$$\begin{aligned} S_1 &\equiv \left\{ s \in I : \frac{1}{n} \sum_{j=1}^n X_j(F(s)) \rightarrow \left[\int X_1 d\nu, - \int -X_1 d\nu \right] \right\}, \\ S_2 &\equiv \left\{ s \in I : \int X_1 d\nu \leq \underline{\lim}_n a_n(\omega) \leq \overline{\lim}_n a_n(\omega) \leq - \int -X_1 d\nu \quad \forall \omega \in F(s) \right\}, \\ \Omega_2 &\equiv \left\{ \omega \in \Omega : \int X_1 d\nu \leq \underline{\lim}_n a_n(\omega) \leq \overline{\lim}_n a_n(\omega) \leq - \int -X_1 d\nu \right\}. \end{aligned}$$

We want to show that $\nu(\Omega_2) = 1$. Notice that

$$\nu(\Omega_2) = \lambda(\{s \in I : F(s) \subseteq \Omega_2\}) = \lambda(S_2).$$

The next Claim will be used to show that $S_1 \subseteq S_2$.

Claim. Let $\{K_n\}$ be a sequence in $\mathcal{K}_{\mathbb{R}}$ such that $K_n \rightarrow [\alpha, \beta]$. Then,

$$\alpha \leq \underline{\lim}_n k_n \leq \overline{\lim}_n k_n \leq \beta$$

for each sequence $\{k_n\}$ in \mathbb{R} such that $k_n \in K_n$ for all $n \geq 1$.

Proof of the Claim: By definition of Hausdorff metric, K_n converges to $[\alpha, \beta]$ if and only if $\max(\max_{t_n \in K_n} \min_{r \in [\alpha, \beta]} |t_n - r|, \max_{r \in [\alpha, \beta]} \min_{t_n \in K_n} |r - t_n|) \rightarrow 0$, in particular

$$\max_{t_n \in K_n} \min_{r \in [\alpha, \beta]} |t_n - r| \rightarrow 0. \quad (1)$$

Let $\{k_{n_j}\}$ be a subsequence of $\{k_n\}$ such that $k_{n_j} \rightarrow \ell \in [-\infty, +\infty]$. If $\ell \notin [\alpha, \beta]$, then there exists $\varepsilon > 0$ such that eventually $|k_{n_j} - r| > \varepsilon$ for all $r \in [\alpha, \beta]$. Hence, we have eventually $\min_{r \in [\alpha, \beta]} |k_{n_j} - r| > \varepsilon$, thus contradicting Eq. (1). \square

If $s \in S_1$, then $\frac{1}{n} \sum_{j=1}^n X_j(F(s)) \rightarrow [\int X_1 d\nu, - \int -X_1 d\nu]$. Hence, for all $\omega \in F(s)$ we have $a_n(\omega) = \frac{1}{n} \sum_{j=1}^n X_j(\omega) \in \frac{1}{n} \sum_{j=1}^n X_j(F(s))$; by the Claim,

$$\int X_1 d\nu \leq \underline{\lim}_n a_n(\omega) \leq \overline{\lim}_n a_n(\omega) \leq - \int -X_1 d\nu.$$

Therefore, $S_1 \subseteq S_2$ and so $\nu(\Omega_2) = \lambda(S_2) \geq \lambda(S_1) = 1$. This completes the proof of the result when (ii) holds.

As to (i), denoting by τ the Polish topology on Ω , there exists a Polish topology $\tau^* \supseteq \tau$ on Ω such that $\sigma(\tau^*) = \mathcal{B}_{\Omega}$, and such that all the X_n s are τ^* -continuous (see, e.g., [21]). Since ν is continuous, then it is a totally monotone capacity with respect to the topology τ^* ; we can thus assume that (ii) holds. \blacksquare

References

- [1] Z. Artstein and R. A. Vitale, “A strong law of large numbers for random compact sets,” *Ann. Probab.* 3, 879-882, 1975.
- [2] R. J. Aumann, “Integrals of set-valued functions,” *J. Math. Anal. Appl.* 12, 1-12, 1965.
- [3] A. Castaldo, F. Maccheroni, and M. Marinacci, “Random sets and their distributions,” mimeo, 2003. <http://web.econ.unito.it/gma/serandom.pdf>
- [4] G. Choquet, “Theory of capacities,” *Ann. Inst. Fourier* 5, 131-295, 1954.
- [5] A. Colubi, M. López-Díaz, J. S. Domínguez-Menchero, and M. A. Gil, “A generalized strong law of large numbers,” *Probab. Theory Relat. Field* 114, 401-417, 1999.
- [6] C. Dellacherie and P-A. Meyer, *Probabilities and potential*, North-Holland, Amsterdam, 1978.
- [7] A. Dempster, “Upper and lower probabilities induced by a multivalued mapping,” *Ann. Math. Statist.* 38, 325-339, 1967.
- [8] J. Doob, *Classical potential theory and its probabilistic counterpart*, Springer-Verlag, Berlin, 1984.
- [9] L. Epstein and M. Schneider, “IID: independently and indistinguishably distributed,” *J. Econom. Theory* 113, 32-50, 2003.
- [10] C. Hess, “Quelques Théorèmes limites pour des ensembles aléatoires bornés ou non,” Séminaire d’Analyse Convexe de l’Université de Montpellier, Exposé 12, 1984.
- [11] P. J. Huber and V. Strassen, “Minimax tests and the Neyman-Pearson lemma for capacities,” *Ann. Statist.* 1, 251-263, 1973.
- [12] E. Klein and A. C. Thompson, *Theory of correspondences*, Wiley, New York, 1984.
- [13] M. Marinacci, “Limit laws for non-additive probabilities and their frequentist interpretation,” *J. Econom. Theory* 84, 145-195, 1999.
- [14] M. Marinacci and L. Montrucchio, “Introduction to the mathematics of ambiguity,” in *Uncertainty in economic theory*, (I. Gilboa, ed.), Routledge, New York, 2004.
- [15] H. T. Nguyen, “On random sets and belief functions,” *J. Math. Anal. Appl.* 65, 531-542, 1978.

- [16] F. Philippe, G. Debs, and J-Y. Jaffray, "Decision making with monotone lower probabilities of infinite order," *Math. Oper. Res.* 24, 767-784, 1999.
- [17] F. N. Proske and M. L. Puri, "A strong law of large numbers for generalized random sets from the viewpoint of empirical processes," *Proc. Amer. Math. Soc.* 131, 2937-2944, 2003.
- [18] D. Schmeidler, "Cores of exact games," *J. Math. Anal. Appl.* 40, 214-225, 1972.
- [19] D. Schmeidler, "Subjective probability and expected utility without additivity," *Econometrica* 57, 571-587, 1989.
- [20] G. Shafer, *A mathematical theory of evidence*, Princeton Univ. Press, Princeton, 1976.
- [21] S. M. Srivastava, *A course on Borel sets*, Springer, New York, 1998.

INTERNATIONAL CENTRE FOR ECONOMIC RESEARCH
APPLIED MATHEMATICS WORKING PAPER SERIES

1. Luigi Montrucchio and Fabio Privileggi, "On Fragility of Bubbles in Equilibrium Asset Pricing Models of Lucas-Type," *Journal of Economic Theory* 101, 158-188, 2001 (ICER WP 2001/5).
2. Massimo Marinacci, "Probabilistic Sophistication and Multiple Priors," *Econometrica* 70, 755-764, 2002 (ICER WP 2001/8).
3. Massimo Marinacci and Luigi Montrucchio, "Subcalculus for Set Functions and Cores of TU Games," *Journal of Mathematical Economics* 39, 1-25, 2003 (ICER WP 2001/9).
4. Juan Dubra, Fabio Maccheroni, and Efe Ok, "Expected Utility Theory without the Completeness Axiom," *Journal of Economic Theory*, forthcoming (ICER WP 2001/11).
5. Adriana Castaldo and Massimo Marinacci, "Random Correspondences as Bundles of Random Variables," April 2001 (ICER WP 2001/12).
6. Paolo Ghirardato, Fabio Maccheroni, Massimo Marinacci, and Marciano Siniscalchi, "A Subjective Spin on Roulette Wheels," *Econometrica*, forthcoming (ICER WP 2001/17).
7. Domenico Menicucci, "Optimal Two-Object Auctions with Synergies," July 2001 (ICER WP 2001/18).
8. Paolo Ghirardato and Massimo Marinacci, "Risk, Ambiguity, and the Separation of Tastes and Beliefs," *Mathematics of Operations Research* 26, 864-890, 2001 (ICER WP 2001/21).
9. Andrea Roncoroni, "Change of Numeraire for Affine Arbitrage Pricing Models Driven By Multifactor Market Point Processes," September 2001 (ICER WP 2001/22).
10. Maitreesh Ghatak, Massimo Morelli, and Tomas Sjöström, "Credit Rationing, Wealth Inequality, and Allocation of Talent", September 2001 (ICER WP 2001/23).
11. Fabio Maccheroni and William H. Ruckle, "BV as a Dual Space," *Rendiconti del Seminario Matematico dell'Università di Padova*, 107, 101-109, 2002 (ICER WP 2001/29).
12. Fabio Maccheroni, "Yaari Dual Theory without the Completeness Axiom," *Economic Theory*, forthcoming (ICER WP 2001/30).
13. Umberto Cherubini and Elisa Luciano, "Multivariate Option Pricing with Copulas," January 2002 (ICER WP 2002/5).
14. Umberto Cherubini and Elisa Luciano, "Pricing Vulnerable Options with Copulas," January 2002 (ICER WP 2002/6).
15. Steven Haberman and Elena Vigna, "Optimal Investment Strategies and Risk Measures in Defined Contribution Pension Schemes," *Insurance: Mathematics and Economics* 31, 35-69, 2002 (ICER WP 2002/10).
16. Enrico Diecidue and Fabio Maccheroni, "Coherence without Additivity," *Journal of Mathematical Psychology*, forthcoming (ICER WP 2002/11).
17. Paolo Ghirardato, Fabio Maccheroni, and Massimo Marinacci, "Ambiguity from the Differential Viewpoint," April 2002 (ICER WP 2002/17).

18. Massimo Marinacci and Luigi Montrucchio, "A Characterization of the Core of Convex Games through Gateaux Derivatives," *Journal of Economic Theory*, forthcoming (ICER WP 2002/18).
19. Fabio Maccheroni and Massimo Marinacci, "How to Cut a Pizza Fairly: Fair Division with Decreasing Marginal Evaluations," *Social Choice and Welfare*, 20, 457-465, 2003 (ICER WP 2002/23).
20. Erio Castagnoli, Fabio Maccheroni and Massimo Marinacci, "Insurance Premia Consistent with the Market," *Insurance: Mathematics and Economics* 31, 267-284, 2002 (ICER WP 2002/24).
21. Fabio Privileggi and Guido Cozzi, "Wealth Polarization and Pulverization in Fractal Societies," September 2002 (ICER WP 2002/39).
22. Paolo Ghirardato, Fabio Maccheroni, and Massimo Marinacci, "Certainty Independence and the Separation of Utility and Beliefs," December 2002 (ICER WP 2002/40).
23. Salvatore Modica and Marco Scarsini, "The Convexity-Cone Approach to Comparative Risk and Downside Risk", January 2003 (ICER WP 2003/1).
24. Claudio Mattalia, "Existence of Solutions and Asset Pricing Bubbles in General Equilibrium Models", January 2003 (ICER WP 2003/2).
25. Massimo Marinacci and Luigi Montrucchio, "Cores and Stable Sets of Finite Dimensional Games", March 2003 (ICER WP 2003/7).
26. Jerome Renault, Sergio Scarlatti, and Marco Scarsini, "A Folk Theorem for Minority Games", April 2003 (ICER WP 2003/10).
27. Peter Klibanoff, Massimo Marinacci, and Sujoy Mukerji, "A Smooth Model of Decision Making under Ambiguity", April 2003 (ICER WP 2003/11).
28. Massimo Marinacci and Luigi Montrucchio, "Ultramodular Functions", June 2003 (ICER WP 2003/13).
29. Erio Castagnoli, Fabio Maccheroni, and Massimo Marinacci, "Choquet Insurance Pricing: a Caveat", June 2003 (ICER WP 2003/14).
30. Thibault Gajdos and Eric Maurin, "Unequal Uncertainties and Uncertain Inequalities: an Axiomatic Approach, June 2003 (ICER WP 2003/15).
31. Thibault Gajdos and John A. Weymark, "Multidimensional Generalized Gini Indices", June 2003 (ICER WP 2003/16).
32. Thibault Gajdos, Jean-Marc Tallon, and Jean-Christophe Vergnaud, "Decision Making with Imprecise Probabilistic Information", June 2003 (ICER WP 2003/18).
33. Alfred Müller and Marco Scarsini, "Archimedean Copulae and Positive Dependence", July 2003 (ICER WP 2003/25).
34. Bruno Bassan, Olivier Gossner, Marco Scarsini, and Shmuel Zamir, "Positive Value of Information in Games", *International Journal of Game Theory*, forthcoming (ICER WP 2003/26).
35. Marco Dall'Aglio and Marco Scarsini, "Zonoids, Linear Dependence, and Size-Biased Distributions on the Simplex", *Advances in Applied Probability* forthcoming (ICER WP 2003/27).
36. Taizhong Hu, Alfred Muller, and Marco Scarsini, "Some Counterexamples in Positive Dependence", *Journal of Statistical Planning and Inference*, forthcoming (ICER WP 2003/28).
37. Massimo Marinacci, Fabio Maccheroni, Alain Chateauneuf, and Jean-Marc Tallon, "Monotone Continuous Multiple Priors" August 2003 (ICER WP 2003/30).

38. Massimo Marinacci and Luigi Montrucchio, “Introduction to the Mathematics of Ambiguity”, October 2003 (ICER WP 2003/34).
39. Fabio Maccheroni, Massimo Marinacci and Aldo Rustichini, “Variational representation of preferences under ambiguity”, June 2004 (ICER WP 2004/5).
40. Antonio Lijoi, Ramsés H. Mena, and Igor Prünster, “Hierarchical mixture modelling with normalized Gaussian priors”, November 2004 (ICER WP 2004/12).
41. Stephen G. Walker, Antonio Lijoi, and Igor Prünster, “Contributions to the understanding of Bayesian consistency”, November 2004 (ICER WP 2004/13).
42. Antonio Lijoi, Igor Prünster, and Stephen G. Walker, “On consistency of nonparametric normal mixtures”, October 2004 (ICER WP 2004/23).
43. Stephen G. Walker, Antonio Lijoi, and Igor Prünster, “On rates of convergence for posterior distributions”, October 2004 (ICER WP 2004/24).
44. Fabio Maccheroni, Massimo Marinacci, Aldo Rustichini and Marco Taboga, “Portfolio selection with monotone mean-variance preferences”, December 2004 (ICER WP 2004/27).
45. Fabio Maccheroni and Massimo Marinacci, “A Strong Law of Large Numbers for Capacities”, December 2004 (ICER WP 2004/28).